

A
Dissertation
on
Acute Hydrocephalus Internus
By Willis Green
of Georgia

Passed March 25th 1823

The human
do unto him
condition, and p
low in the de
most necessarily
to that search a
production of an
time

In closing
school, and a
however, the pr
adaptable con
On oblique to
not in the exp
anxiety, but le
that it may de
land activities
have not been
on their instanc
The subject

The hurried life of a Student of Medicine, who avails himself, of the rich stores of medical education, and practical experience, spread before him in the different lectures of the University, must necessarily preclude the leisure requisite to that research and reflection essential to the production of an original or experimental dissertation

In closing my medical attendance at this school, and ambitious of being clothed with its honours, the preparation of a thesis is rendered an indispensable condition by its regulations.

The obedi^{ance} to this ^{law} precept is this day presented not in the expectation that I can offer any novelty, but limiting my hopes and ambition that it may serve to convince the respected and learned arbiters of my professional capacities that I have not been an idle or inattentive attendant on their instruction

The subject that with this view will be

employed, in the
various, a
below the ana
of his art.

Hydrocephalus
of Physicians.
The early Greek
the time of Galen
to designate some
under the period.

It is however
pathology has
anatomical, the
has been clearly
ability to con
by some writers
Abernethy for
amongst it and
there is no
in other forms

discussed, in these pages is Acute Hydrocephalus
Internus, a disease ^{which} every practitioner has to
deplore the unavailing and inadequate resources
of his art.

Hydrocephalus early attracted the attention
of Physicians. It is noticed in the writings of
the early Greek and Latin authors, though until
the time of Galen, the term was generally applied
to designate serous effusions beneath the skin or
under the pericranium.

It is however only in modern times, since
pathology has been illustrated by post mortem
examinations, that the precise nature of the disease
has been clearly demonstrated, and it may with
propriety be considered as a modern discovery.
But some writers, it has been proposed to remove
Hydrocephalus from the class of dropsies, and to
arrange it among the Phlegmasia or Pyrexia.
There is no solid ground for separating it from
the other forms of dropical affections. The fact

of the analogies
of death;
like the deaths
of the affections
the capillaries of
the brain
are a true mirror
of the state of
the brain.
The proportion
of the capillaries
of the brain
is a true mirror
of the state of
the brain.
The proportion
of the capillaries
of the brain
is a true mirror
of the state of
the brain.

all the analogies of Hydrocephalus, with the other forms of dropsy, prove them to be similar affections. Like the dropsies, too, of the other cavities of the body, the effusion of Hydrocephalus is poured out by the capillaries of a serous membrane, for the Tunica Arachnoides is regarded by modern anatomists as a serous membrane, containing a fluid that lubricates the exterior surface and internal cavity of the brain.

Hydrocephalus may be considered as acute and chronic, idiopathic or sympathetic. The greater proportion of cases that occur are ^{acute} ~~acute~~ ^{sympathetic} ~~acute~~, and proceed from an inflammation, of the Tunica Arachnoides, ^{sympathetically} ~~idiopathically~~ ^{sympathetically} ~~idiopathically~~ affected, and terminating in serous effusion. Acute sympathetic Hydrocephalus, resulting from many different diseases, is the most common, ^{some} in which this affection exists. We find it ensue after some eruptive and other fevers, to be excited by some of the abdominal ^{irritation} ~~disorders~~, by derangement of the digestive organs, and by

the imitation of
Hydrocephalus
in children
the head dentition
it is usually due
to fluid accumulation
or depend to come
has been its
most malady.

Besides the
as to Hydrocephalus
the may increase
but perhaps occ
the head or side
brain, sudden
the suppression of
circulation, the
sudden suppression
in many children
various and

the imitation of intestinal worms.

Hydrocephalus rarely attacks adults, but
sicks on children as its victims. From birth until
the second dentition is the period, during which
it is usually displayed; and then individuals of
a florid complexion, vigorous constitution, subject
or exposed to convulsions, when brothers or sisters
have been its sufferers, are most obnoxious to this
cruel malady.

Besides the causes already stated as giving
rise to Hydrocephalus by Sympathetic imitations
We may enumerate in addition, and as those
that properly occasion it idiosyncratically, blows on
the head or violent concussions, affecting the
brain, sudden terrors, violent paroxysms of anger
the suppression of customary evacuations— as nasal
Hæmorrhage, the matter of crusta Lactea, and the
sudden suppression of transpiration of the head, which,
in many children is very great.

Narcotics and spiritous liquors have been suggested

different countries
of Hydo
known to a
ling; Dr Whitt
abolished State
and it as pro
long, and Rus
of the brain and
Anaphorus, in
Nomenclature pro
as his sign of
some apoplexy
intensity, a depth
thence electro
ing the other co
the insubane a
thousand, occasi
and as imp
tion of these ann
and asphitis

different countries
of Hydo
known to a
ling; Dr Whitt
abolished State
and it as pro
long, and Rus
of the brain and
Anaphorus, in
Nomenclature pro
as his sign of
some apoplexy
intensity, a depth
thence electro
ing the other co
the insubane a
thousand, occasi
and as imp
tion of these ann
and asphitis

by different writers with respect to the primary
cause of Hydrocephalus. Dr Savin attributed
the disease to a torpor of the absorbing vessels of the
brain; Dr Whist considered the effusion to be the result
of a debilitated state of the exhalents; while those who
regarded it as proceeding from inflammation as Bidders
Withering, and Rush, assigned to a general inflamma-
tion of the brain and its membranes, Dr Rush considered
Hydrocephalus, in its first stage, as a less degree of that
inflammation producing phrenitis, and its second,
as a less degree of that effusion which is productive
of serous effusions. I have already stated that modern
anatomists, adopting the doctrine of Richer, class
the *Membræ serosæ* among the serous membranes,
lining the other cavities of the body. Inflammation
of this membrane as inflammation of the Pleura and
Peritoneum, occasions an increase of its serous exhal-
ation, and as inflammation and increased exhal-
ation of these membranes respectively constitute Hydro-
thorax and ascites, so does the inflammation and

permeated whole
substitute the doc
Having thus
to cause, and
I found to a com
treatment.

Acute Rheumatism
attended with
irregular in its
exacerbations and
and violent head
aches or light
intermittent
sweats, with
which is frequent
the virus of the
with a hot burn
to spring to the
head to the head
and left to the

augmented exhalation of the Lunca Arachnoidea,
constitute Hydrocephalus Intermus.

Having thus sketched the history, enumerated
the causes, and stated the nature of Hydrocephalus,
I proceed to a consideration of its symptoms and
Treatment.

Acute Hydrocephalus Hydrocephalus Intermus,
is attended with fever, often not violent, extending
irregular in its paroxysms, having momentary
exacerbations and intermissions; severe constant
and violent headache attends it, exacerbated
by noise or light, and compels the little sufferer
to utter loud screams, deep groans or infant
meanings, with constant complaints of the head
which is frequently in a perpetual state of motion.
The veins of the head are distended, and it imparts
a hot burning sensation to the hand. When
too young to speak, the hands are frequently
brought to the head, but not to any one part of
it, and left to the vacuum than to the face

the, now, and
long as though
something that incre
ely, indeed an
lungs is exhalat
this, as it peg
and irritation of
The pulse part
the poor, sometimes
intermittent, or m
usually towards the
always more fu
hide than the
The tongue is mo
lead; the epigast
nausea and vomit
the gastric affec
long last troubles
the stomach contents
the cross of the dis

eyes, nose, and mouth, with an effort at
picking as though endeavoring to pluck away
something that incommodes. This symptom is gene-
rally held an indication of worms, but it
belongs to cephalic irritation, whether it is sym-
pathic, as it frequently is, from gastric or inter-
stinal irritation or idiosyncrasy.

The pulse partakes of the variable character
of this fever, sometimes frequent and irregular,
intermittent, or natural, or slower than natural,
especially towards the close of the disease, when it
is always more full, and oftentimes more so on
one side than the other.

The tongue is mostly clean or but slightly
furred; the epigastrium region is tender and painful,
nausea and vomiting are rarely absent and
have gastric affections alternate with the headache's
being least troublesome when that is most severe.
Obstinate constipation commonly attends throughout
the course of the disorder, not to be overcome.

have been by
the most as
th a diapha,
and or yellowish
to the air. The
whitish mucous
a liquid and
On the
existence of
at density, the
other character
the discom. The
light with exte
light painful
ly inflamed
When the
are seen to
a mostly but
a brown consi
lets on left

in some cases by the most drastic cathartics. Infants
at the breast are however occasionally affected
with a diarrhea, when the stools are extremely foetid
dark or yellowish, becoming green when exposed
to the air. The urine is scanty, depositing, generally
a whitish mucilaginous matter or which is suspended
in a liquid, under the form of small whitish grains.

On the ^{early} ~~early~~ of symptoms, that announce
the existence of Hydrocephalus, always subject to
great diversity, the eyes of the patient present a
peculiar character, and very distinguishing feature
of the disease. For its commencement, they are
afflicted with extreme sensibility, which renders
the light painful and sometimes they are
highly inflamed.

When the patient is heavy with sleep,
they are seen to roll under the eyelids that
are mostly but half closed. Generally the eyes
are drawn convulsively upwards, so that when
the lids are kept asunder by the fingers, the

under portion of
the disease
on or both eyes
irritation of the
lyst, that has to
acute Hydroce
tentatively to be
late have an
Hydrocephalic
to be remarka
but second to the
and consultations.
belong to intra
statement of intra
At this period
the patient, seem
and incessa
new field
with goudins
dilated pup

under portion of the cornea can alone be seen. As the disease advances Strabismus either of one or both eyes, supervenes, and a peculiar oscillation of the pupil on the admission of light, that has been considered, by some as diagnostic of acute Hydrocephalus, but which is not constant to be met with. The eyes in this state have an expression quite peculiar to the Hydrocephalic Physiology, and which is to be remarked in the intervals of calm that succeed to the painful crises of the head, and convulsions. It is a fixidity that seems to belong to ecstasy, or the expression of profound sentiment of interior contentment.

At this period, the violence of the sufferings of the patient, seem to mitigate; the disturbed sleep and incessant restlessness which existed as first, now yield to a dull and lethargic torpor and with grinding of the teeth, the Strabismus and dilated pupil are increased, and the

has lost
no longer
given to him
in all the
and compass
convulsions
help and a
let quiet
Amphigios
transacted
the face, so
about, are
apophysis
Such
little by ant
Acephalus
develops
It happens
about, and
character,

up has lost its sensibility to the light; admitting no longer distuses the patient, but whatever is given to him is gaudily swallowed. At last arise all the other symptoms of cerebral effusion, and compression. The lethargy becomes more profound, convulsions rack the frame, the breathing is deep and stertorous, the pulse becomes equal but quick and weak scarcely to be counted; Hemiplegia, eyes projecting from their orbits conjunctiva injected, sudden tumefaction of the face, which is partially moistened with sweat, are usually displayed, which is an apoplectic state, which terminates the scene.

Such is the train of formidable formidable symptoms that usually attend on Hydrocephalus Internus. Their progress and development are, however, extremely irregular. It happens at times that most of them are absent, and the disease assumes so insidious a character, that practitioners of great skill

and consider
the true nature
of symptoms of
dyspepsia. After
the alarming
state though
you will appear
in state of health
the symptoms
while paroxysm
become cheerful
among his head
vestige of a
seasonate the
blinded by the
prognostical
light prospect
stew with fix
tation the gate
This moment

and considerable experience have never suspected the true nature of the disease, until the fatal symptoms of the last stage have been suddenly developed. Again, an intermission of all the alarming symptoms will occur, and the patient though touching the last limit of existence will appear to be rapidly returning to a state of health. The stupor will disappear, the symptoms of effusion will vanish, the febrile paroxysms will cease; the infant will become cheerful, take nourishment, and carry his head erect, and no longer present a vestige of disease; the most flattering hopes animate the parents, and the physician deluded by the apparent solution of the disease prognosticates a recovery. But suddenly the bright prospect is overcast, the symptoms of effusion return with frightful rapidity and convulsions hasten the fatal termination of the disease.

This momentary cessation of cerebral congestion

appears to be of
super discharge
dramatic success

The duration
by the great sa
the symptom
the fourth or
the second co
and the dooms

is its course
Treatment. The

patient in the
The first is to
to prevent effusion
when A

To fulfil
had to be
at the same time
to be treated, by
had be warm,

appears to be effected by a profuse salivation, an
copious discharge of semen by Blennorrhoea or an
edematous swelling of the lower extremities. --

The duration of the disease is rendered variable
by the great variation, in the progress, and intensity
of the symptoms. It seldom is protracted beyond
the fourth or terminates before the commencement
of the second week. The more violent the headache
and the sooner the Stomatitis appears the shorter
is its course.

Treatment. Two principle indications govern our
practice in the management of Hydrocephalus Internus.
The first is to abate cerebral irritation, in order
to prevent effusion, and the second is to remove the
effusion when it has taken place.

To fulfil the first indication, measures should
be had to venisection which should be copious,
at the same time local detraction of blood should
be practised, by means of leeches and cups. If the
head be warm, ice should be ^{applied} ~~applied~~ to the head, the hair

being here possible
to immerse in
mustard or
the scalp are
as to maintain
the surface a
ous fluid
Achoer pur
to this purpose
combination with
Clap, cream Tar
superf. Purgative
tly, and should
not advantage,
the stomach, &
administered by the
usual the purg
in double
account of the
this disease.

having been previously removed, and the feet should be immersed in warm water rendered irritating by mustard or pepper. Blisters covering the whole of the scalp are also to be repeatedly applied so as to maintain a constant irritation on the exterior surface and considerable discharge of serous fluid.

Active purging is to be steadily adhered to. For this purpose Calomel, either alone or in combination with ~~the~~^{the} drastic cathartics, as Salap, Enam Tartar, and Gamboge, is to be preferred. Purgative injections are also of great utility, and should not be neglected, they present great advantages, as the extreme irritability of the stomach, causes most of the purgatives, administered by the mouth, to be rejected. In general the purgative medicines should be given in double the doses usually administered on account of the interruption of the primæ viæ in this disease. When however there is great

[Faint, illegible handwriting on the left page, likely bleed-through from the reverse side.]

...and
...purposes
...cases, from
...to be
...The constant
...stopping of
...administration
...ough our
...possessing the
...choice to be
...employment
...Hypnotism
...are said
...The second
...Hypnotism
...it has
...disputed
...has been
...with great
...in the

hardness and tenderness of the abdomen, the powerful purgatives may prove injurious; and in these cases, fomentations and a semicupium are to be promised.

The constant nausea and vomiting are distressing symptoms, that interfere with the administration of Medicines, and require attention; though our remedies are seldom successful in removing them. Dr Fothergill advises *Emeticon Rubiacum* to be given with this view, but its employment we believe to be injurious, Antispasmodics applied externally, to the epigastric region are said to be sometimes successful.

The second indication in the treatment of *Hydrocephalus* is to dissipate the effusion when it has occurred, and which may be pursued simultaneously with the first. Mercury has been advised by Drs Percival and Lobson, with great confidence, administered internally and in the form of unctuous, so as to affect

the salary given
the great success
of these gentlemen
the assistant of
the preparations
with the admini-
strative efforts
Vothius has
constant discha-
of Africa and
to this purpose.
The preceding
we had suited
Hobbes & others;
the of the disci-
with a prospect,
we shall have
abstract form. a

the salivary glands. Experience has not justified the great encomiums, passed on this medicine by those gentlemen, but from its power of exciting the absorbent system, it should not be omitted.

The preparations of Squills should be combined with the administration of Mercury, to produce a diuretic effect or action.

Cochineus have been advised to keep up a constant discharge from the nostrils. The flowers of Snice and Macaba snuff have been recommended for this purpose.

The preceding methodus medendi appears to me best suited to the treatment of Acute Syphilis - Pholis Intestina; but it is only in the forming stage of the disease that we can flatter ourselves with a prospect, by pursuing it with vigour, that we shall have the satisfaction of rescuing our patient from a premature grave.

Inauguration

Hydro

Alex